

JORDAN DERIVATIONS OF INCIDENCE ALGEBRAS

ZHANKUI XIAO

ABSTRACT. Let \mathcal{R} be a commutative ring with identity, $I(X, \mathcal{R})$ be the incidence algebra of a locally finite pre-ordered set X . In this note, we characterise the derivations of $I(X, \mathcal{R})$ and prove that every Jordan derivation of $I(X, \mathcal{R})$ is a derivation provided that \mathcal{R} is 2-torsion free.

1. INTRODUCTION

Let \mathcal{R} be a commutative ring with identity, A be an algebra over \mathcal{R} . An \mathcal{R} -linear mapping $D : A \rightarrow A$ is called a derivation if $D(xy) = D(x)y + xD(y)$ for all $x, y \in A$, and is called a *Jordan derivation* if

$$D(x^2) = D(x)x + xD(x)$$

for all $x \in A$. There has been a great interest in the study of Jordan derivations of various algebras in the last decades. The standard problem is to find out whether a Jordan derivation degenerate to a derivation. Jacobson and Rickart [8] proved that every Jordan derivation of the full matrix algebra over a 2-torsion free unital ring is a derivation by relating the problem to the decomposition of Jordan homomorphisms. In [7], Herstein showed that every Jordan derivation from a 2-torsion free prime ring into itself is also a derivation. These results have been extended to different rings and algebras in various directions (see [2, 3, 6, 11, 20] and the references therein). We would like to refer the reader to Brešar's paper [4] for a comprehensive and more detailed understanding of this topic.

We now recall the definition of incidence algebras. Let (X, \leq) be a locally finite pre-ordered set. This means \leq is a reflexive and transitive binary relation on the set X , and for any $x \leq y$ in X there are only finitely many elements z satisfying $x \leq z \leq y$. The *incidence algebra* $I(X, \mathcal{R})$ of X over \mathcal{R} is defined on the set

$$I(X, \mathcal{R}) := \{f : X \times X \longrightarrow \mathcal{R} \mid f(x, y) = 0 \text{ if } x \not\leq y\}$$

with algebraic operation given by

$$\begin{aligned} (f + g)(x, y) &= f(x, y) + g(x, y), \\ (rf)(x, y) &= rf(x, y), \\ (fg)(x, y) &= \sum_{x \leq z \leq y} f(x, z)g(z, y) \end{aligned}$$

for all $f, g \in I(X, \mathcal{R})$, $r \in \mathcal{R}$ and $x, y, z \in X$. The product (fg) is usually called *convolution* in function theory. It would be helpful to point out that the full matrix

2010 *Mathematics Subject Classification.* 16W10, 16W25, 47L35.

Key words and phrases. derivation, Jordan derivation, incidence algebra.

The work was supported by a research foundation of NSFC. (Grant No. 11226068).

algebra $M_n(\mathcal{R})$ and the upper (or lower) triangular matrix algebras $T_n(\mathcal{R})$ are special examples of incidence algebras

Incidence algebras were first considered by Ward [18] as generalized algebras of arithmetic functions. Rota and Stanley developed incidence algebras as the fundamental structures of enumerative combinatorial theory and allied areas of arithmetic function theory (see [15]). Motivated by the results of Stanley [17], automorphisms and other algebraic mappings of incidence algebras have been extensively studied (see [1, 5, 9, 10, 12, 13, 14, 16] and the references therein). Baclawski [1] studied the automorphisms and derivations of incidence algebras $I(X, \mathcal{R})$ when X is a locally finite partially ordered set. Especially, he proved that every derivation of $I(X, \mathcal{R})$ with X a locally finite partially ordered set can be decomposed as a sum of an inner derivation and a transitive induced derivation (see the definition in Section 2). Koppinen [10] has extended these results to the incidence algebras $I(X, \mathcal{R})$ with X a locally finite pre-ordered set. In present paper, we first characterise the derivations of $I(X, \mathcal{R})$ by a direct computation. Based on such characterization of derivations, we prove that every Jordan derivation of $I(X, \mathcal{R})$ is a derivation provided that \mathcal{R} is 2-torsion free.

2. DERIVATIONS ON INCIDENCE ALGEBRAS

In this section, we characterise the derivations of the incidence algebras $I(X, \mathcal{R})$ where X is a locally finite pre-ordered set. We first single out some properties of $I(X, \mathcal{R})$ for later use. One can find out all information in [15] and [10].

The identity element δ of $I(X, \mathcal{R})$ is given by $\delta(x, y) = \delta_{xy}$ for $x \leq y$, where $\delta_{xy} \in \{0, 1\}$ is the Kronecker delta. If $x, y \in X$ with $x \leq y$, let e_{xy} be defined by $e_{xy}(u, v) = 1$ if $(u, v) = (x, y)$, and $e_{xy}(u, v) = 0$ otherwise. Then $e_{xy}e_{uv} = \delta_{yu}e_{xv}$ by the definition of convolution. Moreover, the set $\mathfrak{B} := \{e_{xy} \mid x \leq y\}$ forms an \mathcal{R} -linear basis of $I(X, \mathcal{R})$. The following lemma, to some extent, is well-known and we give out the proof here for completeness.

Lemma 2.1. *Let A be an \mathcal{R} -algebra with an \mathcal{R} -linear basis Y . Then an \mathcal{R} -linear operator $D : A \rightarrow A$ is a derivation if and only if*

$$D(xy) = D(x)y + xD(y), \quad \forall x, y \in Y.$$

Proof. We only need to prove the sufficiency. Let $a, b \in A$ and

$$a = \sum_{x \in Y} C_x x, \quad b = \sum_{y \in Y} C'_y y,$$

where $C_x, C'_y \in \mathcal{R}$. Then

$$\begin{aligned} D(ab) &= D\left(\sum_{x, y \in Y} C_x C'_y xy\right) \\ &= \sum_{x, y \in Y} C_x C'_y (D(x)y + xD(y)) \\ &= D\left(\sum_{x \in Y} C_x x\right) \left(\sum_{y \in Y} C'_y y\right) + \left(\sum_{x \in Y} C_x x\right) D\left(\sum_{y \in Y} C'_y y\right) \\ &= D(a)b + aD(b). \end{aligned}$$

□

Let $D : I(X, \mathcal{R}) \rightarrow I(X, \mathcal{R})$ be an \mathcal{R} -linear operator. Motivated by the above Lemma, we denote for all $i, j \in X$ with $i \leq j$

$$D(e_{ij}) = \sum_{e_{xy} \in \mathfrak{B}} C_{xy}^{ij} e_{xy}.$$

We will use this notation for the rest of this paper. For any $x \in X$, define L_x and R_x to be the subsets of X by

$$L_x := \{i \in X \mid i \leq x, i \neq x\} \quad \text{and} \quad R_x := \{j \in X \mid x \leq j, j \neq x\}.$$

Note that the set $L_x \cap R_x$ maybe not empty since X is a pre-ordered set.

Theorem 2.2. *Let $D : I(X, \mathcal{R}) \rightarrow I(X, \mathcal{R})$ be an \mathcal{R} -linear operator. Then D is a derivation if and only if D satisfies*

$$D(e_{ij}) = \sum_{x \in L_i} C_{xi}^{ii} e_{xj} + C_{ij}^{ij} e_{ij} + \sum_{y \in R_j} C_{jy}^{jj} e_{iy} \quad (1)$$

for all $e_{ij} \in \mathfrak{B}$, where the coefficients C_{xy}^{ij} are subject to the following relations

$$\begin{aligned} C_{jk}^{jj} + C_{jk}^{kk} &= 0, \quad \text{if } j \leq k; \\ C_{ij}^{ij} + C_{jk}^{jk} &= C_{ik}^{ik}, \quad \text{if } i \leq j, j \leq k. \end{aligned}$$

Proof. (\Rightarrow). For any $i \in X$, there is $D(e_{ii}) = D(e_{ii})e_{ii} + e_{ii}D(e_{ii})$. Thus

$$\begin{aligned} \sum_{e_{xy} \in \mathfrak{B}} C_{xy}^{ii} e_{xy} &= \left(\sum_{e_{xy} \in \mathfrak{B}} C_{xy}^{ii} e_{xy} \right) e_{ii} + e_{ii} \left(\sum_{e_{xy} \in \mathfrak{B}} C_{xy}^{ii} e_{xy} \right) \\ &= \sum_{x, x \leq i} C_{xi}^{ii} e_{xi} + \sum_{y, i \leq y} C_{iy}^{ii} e_{iy}. \end{aligned} \quad (2)$$

Taking $x = i$ and $y = i$ in (2), we have $C_{ii}^{ii} = 0$. Hence

$$D(e_{ii}) = \sum_{x \in L_i} C_{xi}^{ii} e_{xi} + \sum_{y \in R_i} C_{iy}^{ii} e_{iy}. \quad (3)$$

For any $e_{ij} \in \mathfrak{B}$ with $i \neq j$, we deduce from equation (3) that

$$\begin{aligned} D(e_{ij}) &= D(e_{ii}e_{ij}e_{jj}) \\ &= D(e_{ii})e_{ij} + e_{ii}D(e_{ij})e_{jj} + e_{ij}D(e_{jj}) \\ &= D(e_{ii})e_{ij} + C_{ij}^{ij} e_{ij} + e_{ij}D(e_{jj}) \\ &= \sum_{x \in L_i} C_{xi}^{ii} e_{xj} + C_{ij}^{ij} e_{ij} + \sum_{y \in R_j} C_{jy}^{jj} e_{iy}. \end{aligned} \quad (4)$$

Note that the identities (3) and (4) proves (1).

In order to determine the coefficients C_{xy}^{ij} , we need to apply the derivation D to the identity $e_{ij}e_{kl} = \delta_{jk}e_{il}$. There are two subcases.

Case 1. When $j \neq k$, then

$$\begin{aligned} 0 &= D(e_{ij})e_{kl} + e_{ij}D(e_{kl}) \\ &= \left(\sum_{x \in L_i} C_{xi}^{ii} e_{xj} + C_{ij}^{ij} e_{ij} + \sum_{y \in R_j} C_{jy}^{jj} e_{iy} \right) e_{kl} \\ &\quad + e_{ij} \left(\sum_{x \in L_k} C_{xk}^{kk} e_{xl} + C_{kl}^{kl} e_{kl} + \sum_{y \in R_l} C_{ly}^{ll} e_{ky} \right) \\ &= C_{jk}^{jj} e_{il} + C_{jk}^{kk} e_{il}. \end{aligned} \quad (5)$$

Therefore $C_{jk}^{jj} + C_{jk}^{kk} = 0$.

Case 2. When $j = k$, then

$$\begin{aligned}
D(e_{il}) &= D(e_{ij})e_{jl} + e_{ij}D(e_{jl}) \\
&= \left(\sum_{x \in L_i} C_{xi}^{ii}e_{xj} + C_{ij}^{ij}e_{ij} + \sum_{y \in R_j} C_{jy}^{jj}e_{iy} \right) e_{jl} \\
&\quad + e_{ij} \left(\sum_{x \in L_j} C_{xj}^{jj}e_{xl} + C_{jl}^{jl}e_{jl} + \sum_{y \in R_l} C_{ly}^{ll}e_{jy} \right) \\
&= \sum_{x \in L_i} C_{xi}^{ii}e_{xl} + C_{ij}^{ij}e_{il} + C_{jl}^{jl}e_{il} + \sum_{y \in R_l} C_{ly}^{ll}e_{iy}.
\end{aligned} \tag{6}$$

Taking (1) into account, we obtain $C_{ij}^{ij} + C_{jl}^{jl} = C_{il}^{il}$, which proves the necessity.

(\Leftarrow). First it follows from the equation $C_{ij}^{ij} + C_{jk}^{jk} = C_{ik}^{ik}$ that $C_{ii}^{ii} = 0$ for all $i \in X$. Hence by (1), $D(e_{ii})$ is of the form as in equation (3), i.e.

$$D(e_{ii}) = \sum_{x \in L_i} C_{xi}^{ii}e_{xi} + \sum_{y \in R_i} C_{iy}^{ii}e_{iy}. \tag{7}$$

Now by Lemma 2.1, to show that D is a derivation we just need to show that $D(e_{ij}e_{kl}) = D(e_{ij})e_{kl} + e_{ij}D(e_{kl})$ for all $e_{ij}, e_{kl} \in \mathfrak{B}$. On the other hand, a little modification of the identities (5) and (6) shows that the equations $D(e_{ij}e_{kl}) = D(e_{ij})e_{kl} + e_{ij}D(e_{kl})$ for all $e_{ij}, e_{kl} \in \mathfrak{B}$ always hold. This completes the proof of the theorem. \square

For any $g \in I(X, \mathcal{R})$, the mapping $f \mapsto [g, f]$ is a derivation of $I(X, \mathcal{R})$. We call it an *inner derivation* and denote it as Inn_g . A mapping $f : \leq \rightarrow \mathcal{R}$ is called *transitive* if

$$f(i, j) + f(j, k) = f(i, k)$$

for all $i, j, k \in X$ such that $i \leq j, j \leq k$. Note that every mapping $\sigma : X \rightarrow \mathcal{R}$ determines a transitive mapping $(i, j) \mapsto \sigma(i) - \sigma(j)$. Transitive mappings of this form are called *trivial*. If $f : \leq \rightarrow \mathcal{R}$ is a transitive mapping, we define an \mathcal{R} -linear mapping $\Delta_f : I(X, \mathcal{R}) \rightarrow I(X, \mathcal{R})$ by

$$\Delta_f(e_{ij}) = f(i, j)e_{ij}$$

for all $e_{ij} \in \mathfrak{B}$.

Lemma 2.3. *With notations as above, Δ_f is a derivation. Moreover Δ_f is inner if and only if f is trivial.*

Proof. Note that $\Delta_f(e_{ij})e_{kl} + e_{ij}\Delta_f(e_{kl}) = \delta_{jk}(f(i, j) + f(k, l))e_{il} = \Delta_f(e_{ij}e_{kl})$ for all $e_{ij}, e_{kl} \in \mathfrak{B}$. Then Lemma 2.1 implies that Δ_f is a derivation.

If f is trivial, there exists a mapping $\sigma : X \rightarrow \mathcal{R}$ such that $f(i, j) = \sigma(i) - \sigma(j)$. Let $g = \sum_{i \in X} \sigma(i)e_{ii} \in I(X, \mathcal{R})$. Then $[g, e_{ij}] = (\sigma(i) - \sigma(j))e_{ij} = \Delta_f(e_{ij})$ for all $e_{ij} \in \mathfrak{B}$, and hence Δ_f is inner. Conversely, if Δ_f is inner, there exists an element $g = \sum_{e_{ij} \in \mathfrak{B}} C_{ij}e_{ij}$ such that $\Delta_f(e_{ij}) = [g, e_{ij}]$ for all $e_{ij} \in \mathfrak{B}$. Hence $f(i, j) = C_{ii} - C_{jj}$. Setting $\sigma : X \rightarrow \mathcal{R}$ by $\sigma(i) = C_{ii}$, we get that f is trivial. \square

Remark 2.4. We will call the derivation Δ_f by *transitive induced derivation* induced by the transitive mapping f . The definition of transitive mappings first appeared in Nowicki's paper [13], where he dealt with finite incidence algebras, i.e. X is a finite pre-ordered set.

Proposition 2.5. *Let D be a derivation of $I(X, \mathcal{R})$. Then there exist an element $g \in I(X, \mathcal{R})$ and a transitive mapping f such that*

$$D = \text{Inn}_g + \Delta_f.$$

Proof. It follows from Theorem 2.2 that

$$D(e_{ij}) = \sum_{x \in L_i} C_{xi}^{ii} e_{xj} + C_{ij}^{ij} e_{ij} + \sum_{y \in R_j} C_{jy}^{jj} e_{iy}$$

for all $e_{ij} \in \mathfrak{B}$, where the coefficients C_{xy}^{ij} are subject to the following relations

$$C_{jk}^{jj} + C_{jk}^{kk} = 0 \quad \text{and} \quad C_{ij}^{ij} + C_{jk}^{jk} = C_{ik}^{ik}.$$

Define $f : \leq \rightarrow \mathcal{R}$ by $(i, j) \mapsto C_{ij}^{ij}$. Then f is transitive since $C_{ij}^{ij} + C_{jk}^{jk} = C_{ik}^{ik}$. On the other hand, Let $g = \sum_{e_{ij} \in \mathfrak{B}} C_{ij}^{jj} e_{ij}$. Note that $g(i, j) = C_{ij}^{jj} = -C_{ij}^{ii}$. Hence

$$D(e_{ij}) = \text{Inn}_g(e_{ij}) + \Delta_f(e_{ij})$$

for all $e_{ij} \in \mathfrak{B}$. The proposition follows from the fact that the mappings D , Inn_g and Δ_f are all \mathcal{R} -linear. \square

We point out that Proposition 2.5 can also be obtained from [10, Theorem 6.3], where Koppinen use deeply the coalgebra structure of incidence algebras. Our approach is elementary but is convenient for studying Jordan derivations in next section. Lemma 2.3 and Proposition 2.5 deduce the following.

Corollary 2.6. *Every derivation of $I(X, \mathcal{R})$ is inner if and only if every transitive mapping is trivial.*

Corollary 2.7. *The incidence algebra $I(X, \mathcal{R})$ is a Kadison algebra, i.e. every local derivation of $I(X, \mathcal{R})$ is a derivation.*

Proof. The proof is similar to that of [14, Theorem 3]. \square

3. JORDAN DERIVATIONS ON INCIDENCE ALGEBRAS

Throughout this section, we assume that \mathcal{R} is 2-torsion free. Let $D : I(X, \mathcal{R}) \rightarrow I(X, \mathcal{R})$ be a Jordan derivation. We also denote

$$D(e_{ij}) = \sum_{e_{xy} \in \mathfrak{B}} C_{xy}^{ij} e_{xy}.$$

for all $i, j \in X$ with $i \leq j$. We assume $C_{xy}^{ij} = 0$, if needed, for $x \not\leq y$. The following lemma is due to Herstein [7].

Lemma 3.1. *Let A be a 2-torsion free ring, $D : A \rightarrow A$ be a Jordan derivation. Then for any $a, b, c \in A$, we have*

- (1) $D(ab + ba) = D(a)b + aD(b) + D(b)a + bD(a)$,
- (2) $D(aba) = D(a)ba + aD(b)a + abD(a)$,
- (3) $D(abc + cba) = D(a)bc + aD(b)c + abD(c) + D(c)ba + cD(b)a + cbD(a)$.

Lemma 3.2. *Let $D : I(X, \mathcal{R}) \rightarrow I(X, \mathcal{R})$ be a Jordan derivation. Then*

$$D(e_{ij}) = \sum_{x \in L_i} C_{xi}^{ii} e_{xj} + C_{ij}^{ij} e_{ij} + \sum_{y \in R_j} C_{jy}^{jj} e_{iy} + C_{ji}^{ij} e_{ji}$$

for all $e_{ij} \in \mathfrak{B}$, where the coefficients C_{xy}^{ij} are subject to the following relations

$$\begin{aligned} C_{jk}^{jj} + C_{jk}^{kk} &= 0, \quad \text{if } j \leq k; \\ C_{ij}^{ij} + C_{jk}^{jk} &= C_{ik}^{ik}, \quad \text{if } i \leq j, j \leq k. \end{aligned}$$

Proof. Since e_{ii} is an idempotent element for all $i \in X$, a similar computation as in Theorem 2.2 shows that $C_{ii}^{ii} = 0$ and

$$D(e_{ii}) = \sum_{x \in L_i} C_{xi}^{ii} e_{xi} + \sum_{y \in R_i} C_{iy}^{ii} e_{iy}. \quad (8)$$

For any $e_{ij} \in \mathfrak{B}$ with $i \neq j$, we deduce from Lemma 3.1 (3) and equation (8) that

$$\begin{aligned} D(e_{ij}) &= D(e_{ii}e_{ij}e_{jj} + e_{jj}e_{ij}e_{ii}) \\ &= D(e_{ii})e_{ij} + e_{ii}D(e_{ij})e_{jj} + e_{ij}D(e_{jj}) + e_{jj}D(e_{ij})e_{ii} \\ &= D(e_{ii})e_{ij} + C_{ij}^{ij}e_{ij} + e_{ij}D(e_{jj}) + C_{ji}^{ij}e_{ji} \\ &= \sum_{x \in L_i} C_{xi}^{ii}e_{xj} + C_{ij}^{ij}e_{ij} + \sum_{y \in R_j} C_{jy}^{jj}e_{iy} + C_{ji}^{ij}e_{ji}. \end{aligned} \quad (9)$$

Combining the equations (8) and (9), we get the form of $D(e_{ij})$. It would be helpful to point out that $C_{ji}^{ij} = 0$ if $j \not\leq i$ by our assumption.

In order to determine the coefficients C_{xy}^{ij} , we need to apply the Jordan derivation D to the identity $e_{ij}e_{kl} + e_{kl}e_{ij} = \delta_{jk}e_{il} + \delta_{li}e_{kj}$. There are four subcases.

Case 1. When $l \neq i$ and $j \neq k$, we have

$$\begin{aligned} 0 &= D(e_{ij})e_{kl} + e_{ij}D(e_{kl}) + D(e_{kl})e_{ij} + e_{kl}D(e_{ij}) \\ &= \left(\sum_{x \in L_i} C_{xi}^{ii}e_{xj} + C_{ij}^{ij}e_{ij} + \sum_{y \in R_j} C_{jy}^{jj}e_{iy} + C_{ji}^{ij}e_{ji} \right) e_{kl} \\ &\quad + e_{ij} \left(\sum_{x \in L_k} C_{xk}^{kk}e_{xl} + C_{kl}^{kl}e_{kl} + \sum_{y \in R_l} C_{ly}^{ll}e_{ky} + C_{lk}^{kl}e_{lk} \right) \\ &\quad + \left(\sum_{x \in L_k} C_{xk}^{kk}e_{xl} + C_{kl}^{kl}e_{kl} + \sum_{y \in R_l} C_{ly}^{ll}e_{ky} + C_{lk}^{kl}e_{lk} \right) e_{ij} \\ &\quad + e_{kl} \left(\sum_{x \in L_i} C_{xi}^{ii}e_{xj} + C_{ij}^{ij}e_{ij} + \sum_{y \in R_j} C_{jy}^{jj}e_{iy} + C_{ji}^{ij}e_{ji} \right) \\ &= C_{jk}^{jj}e_{il} + \delta_{ik}C_{ji}^{ij}e_{jl} + C_{jk}^{kk}e_{il} + \delta_{jl}C_{lk}^{kl}e_{ik} \\ &\quad + C_{li}^{ll}e_{kj} + \delta_{ik}C_{lk}^{kl}e_{lj} + C_{li}^{ii}e_{kj} + \delta_{jl}C_{ji}^{ij}e_{ki}. \end{aligned} \quad (10)$$

We consider the coefficient of e_{il} in the equation (10). If $e_{il} = e_{jl}$, then $C_{ji}^{ij} = C_{ii}^{ii} = 0$. If $e_{il} = e_{ik}$, then $C_{lk}^{kl} = C_{kk}^{kk} = 0$. Therefore, $C_{jk}^{jj} + C_{jk}^{kk} = 0$ if $(i, l) \neq (k, j)$ or $2(C_{jk}^{jj} + C_{jk}^{kk}) = 0$ if $(i, l) = (k, j)$. Hence $C_{jk}^{jj} + C_{jk}^{kk} = 0$ for $j \leq k$ since \mathcal{R} is 2-torsion free.

Case 2. When $l \neq i$ and $j = k$, we have

$$\begin{aligned}
D(e_{il}) &= D(e_{ij})e_{kl} + e_{ij}D(e_{kl}) + D(e_{kl})e_{ij} + e_{kl}D(e_{ij}) \\
&= \left(\sum_{x \in L_i} C_{xi}^{ii}e_{xj} + C_{ij}^{ij}e_{ij} + \sum_{y \in R_j} C_{jy}^{jj}e_{iy} + C_{ji}^{ij}e_{ji} \right) e_{jl} \\
&\quad + e_{ij} \left(\sum_{x \in L_j} C_{xj}^{jj}e_{xl} + C_{jl}^{jl}e_{jl} + \sum_{y \in R_l} C_{ly}^{ll}e_{jy} + C_{lj}^{jl}e_{lj} \right) \\
&\quad + \left(\sum_{x \in L_j} C_{xj}^{jj}e_{xl} + C_{jl}^{jl}e_{jl} + \sum_{y \in R_l} C_{ly}^{ll}e_{jy} + C_{lj}^{jl}e_{lj} \right) e_{ij} \\
&\quad + e_{jl} \left(\sum_{x \in L_i} C_{xi}^{ii}e_{xj} + C_{ij}^{ij}e_{ij} + \sum_{y \in R_j} C_{jy}^{jj}e_{iy} + C_{ji}^{ij}e_{ji} \right) \\
&= \sum_{x \in L_i} C_{xi}^{ii}e_{xl} + C_{ij}^{ij}e_{il} + \delta_{ij}C_{ji}^{ij}e_{jl} + C_{jl}^{jl}e_{il} + \sum_{y \in R_l} C_{ly}^{ll}e_{iy} + \delta_{jl}C_{lj}^{jl}e_{ij} \\
&\quad + C_{li}^{ll}e_{jj} + \delta_{ij}C_{lj}^{jl}e_{lj} + C_{li}^{ii}e_{jj} + \delta_{lj}C_{ji}^{ij}e_{ji} \\
&= \sum_{x \in L_i} C_{xi}^{ii}e_{xl} + C_{ij}^{ij}e_{il} + C_{jl}^{jl}e_{il} + \sum_{y \in R_l} C_{ly}^{ll}e_{iy} + \delta_{ij}C_{lj}^{jl}e_{lj} + \delta_{lj}C_{ji}^{ij}e_{ji},
\end{aligned} \tag{11}$$

where the last identity follows from the fact $\delta_{ij}C_{ji}^{ij} = 0$ and $C_{li}^{ll} + C_{li}^{ii} = 0$. Therefore, there is

$$C_{il}^{il}e_{il} + C_{li}^{il}e_{li} = C_{ij}^{ij}e_{il} + C_{jl}^{jl}e_{il} + \delta_{ij}C_{lj}^{jl}e_{lj} + \delta_{lj}C_{ji}^{ij}e_{ji}. \tag{12}$$

We consider the coefficient of e_{il} in the equation (12). Note that $l \neq i$ in this case. We obtain $C_{ij}^{ij} + C_{jl}^{jl} = C_{il}^{il}$.

Case 3. When $l = i$ and $j \neq k$, this case is same with the Case 2.

Case 4. When $l = i$ and $j = k$, a direct calculation shows that $C_{ij}^{ij} + C_{ji}^{ji} = 0$. Combining with the Case 2 or the Case 3, we have $C_{ij}^{ij} + C_{jl}^{jl} = C_{il}^{il}$ for all $i \leq j$ and $j \leq l$. \square

We now are at the position to prove the main result of this paper.

Theorem 3.3. *Let \mathcal{R} be a 2-torsion free commutative ring with identity. Then every Jordan derivation of the incidence algebra $I(X, \mathcal{R})$ is a derivation.*

Proof. Let $D : I(X, \mathcal{R}) \rightarrow I(X, \mathcal{R})$ be a Jordan derivation. By Lemma 3.2

$$D(e_{ij}) = \sum_{x \in L_i} C_{xi}^{ii}e_{xj} + C_{ij}^{ij}e_{ij} + \sum_{y \in R_j} C_{jy}^{jj}e_{iy} + C_{ji}^{ij}e_{ji}$$

for all $e_{ij} \in \mathfrak{B}$, where the coefficients C_{xy}^{ij} are subject to the following relations

$$\begin{aligned}
C_{jk}^{jj} + C_{jk}^{kk} &= 0, \quad \text{if } j \leq k; \\
C_{ij}^{ij} + C_{jk}^{jk} &= C_{ik}^{ik}, \quad \text{if } i \leq j, j \leq k.
\end{aligned}$$

We define an \mathcal{R} -linear operator d by

$$d(e_{ij}) = \sum_{x \in L_i} C_{xi}^{ii}e_{xj} + C_{ij}^{ij}e_{ij} + \sum_{y \in R_j} C_{jy}^{jj}e_{iy}$$

for all $e_{ij} \in \mathfrak{B}$, where the coefficients C_{xy}^{ij} are subject to the following relations

$$\begin{aligned} C_{jk}^{jj} + C_{jk}^{kk} &= 0, \text{ if } j \leq k; \\ C_{ij}^{ij} + C_{jk}^{jk} &= C_{ik}^{ik}, \text{ if } i \leq j, j \leq k. \end{aligned}$$

Then Theorem 2.2 makes d be a derivation. Then the operator $\Delta := D - d$ is a Jordan derivation of $I(X, \mathcal{R})$ satisfying

$$\Delta(e_{ii}) = 0 \text{ and } \Delta(e_{ij}) = C_{ji}^{ij} e_{ji}$$

for all $i \leq j$. Note that we have assumed $C_{ji}^{xy} = 0$ if $j \not\leq i$. Restrict Δ to the subalgebra generated by $e_{ii}, e_{jj}, e_{ij}, e_{ji}$ if $j \leq i$ which is isomorphic to the full matrix algebra $M_2(\mathcal{R})$. Then [8, Theorem 7 and Theorem 22] imply that Δ is a derivation and hence $C_{ji}^{ij} = 0$. This completes the proof of the theorem. \square

Remark 3.4. If X is a finite partially ordered set (poset), then the incidence algebra $I(X, \mathcal{R})$ (also called bigraph algebra or finite dimensional CSL algebra) is usually a triangular algebra (see [19, P.1245]). At this case, Theorem 3.3 can be deduced from [11, Theorem 3.2] or [20, Theorem 2.1].

REFERENCES

- [1] K. Baclawski, *Automorphisms and derivations of incidence algebras*, Proc. Amer. Math. Soc., **36** (1972), 351-356.
- [2] M. Brešar, *Jordan derivations on semiprime rings*, Proc. Amer. Math. Soc., **104** (1988), 1003-1006.
- [3] M. Brešar, *Jordan mappings of semiprime rings*, J. Algebra, **127** (1989), 218-228.
- [4] M. Brešar, *Jordan derivations revisited*, Math. Proc. Camb. Phil. Soc., **139** (2005), 411-425.
- [5] S. P. Coelho and C. P. Milies, *Derivations of upper triangular matrix rings*, Linear Algebra Appl., **187** (1993), 263-267.
- [6] J. M. Cusack, *Jordan derivations on rings*, Proc. Amer. Math. Soc., **53** (1975), 321-324.
- [7] I. N. Herstein, *Jordan derivations of prime rings*, Proc. Amer. Math. Soc., **8** (1957), 1104-1110.
- [8] N. Jacobson and C. Rickart, *Jordan homomorphisms of rings*, Trans. Amer. Math. Soc., **69** (1950), 479-502.
- [9] S. Jøndrup, *Automorphisms and derivations of upper triangular matrix rings*, Linear Algebra Appl., **221** (1995), 205-218.
- [10] M. Koppinen, *Automorphisms and higher derivations of incidence algebras*, J. Algebra., **174** (1995), 698-723.
- [11] F. Lu, *The Jordan structure of CSL algebras*, Studia Math., **190** (2009), 283-299.
- [12] D. Mathis, *Differential polynomial rings and Morita equivalence*, Comm. Algebra, **10** (1982), 2001-2017.
- [13] A. Nowicki, *Derivations of special subrings of matrix rings and regular graphs*, Tsukuba J. Math., **7** (1983), 281-297.
- [14] A. Nowicki and I. Nowosad, *Local derivations of subrings of matrix rings*, Acta Math. Hungar., **105** (2004), 145-150.
- [15] E. Spiegel and C. O'Donnell, *Incidence algebras*, Monographs and Textbooks in Pure and Applied Mathematics, vol. **206**, marcel Dekker, New York, 1997.
- [16] E. Spiegel, *On the automorphisms of incidence algebras*, J. Algebra, **239** (2001), 615-623.
- [17] R. Stanley, *Structure of incidence algebras and their automorphism groups*, Bull. Amer. Math. Soc., **76** (1970), 1236-1239.
- [18] M. Ward, *Arithmetic functions on rings*, Ann. Math., **38** (1937), 725-732.
- [19] Z.-K. Xiao and F. Wei, *Lie triple derivations of triangular algebras*, Linear Algebra Appl., **437** (2012), 1234-1249.
- [20] J.-H. Zhang and W.-Y. Yu, *Jordan derivations of triangular algebras*, Linear Algebra Appl., **419** (2006), 251-255.

XIAO: SCHOOL OF MATHEMATICAL SCIENCES, HUAQIAO UNIVERSITY, QUANZHOU, FUJIAN, 362021,
P. R. CHINA
E-mail address: zhkxiao@gmail.com